






The ER Protein Translocation Channel Subunit Sbh1 Controls Virulence of *Cryptococcus neoformans*

 Felipe H. Santiago-Tirado,^{a*} Thomas Hurtaux,^a  Jennifer Geddes-McAlister,^b Duy Nguyen,^{c§} Volkhard Helms,^c
 Tamara L. Doering,^a Karin Römisch^d

^aDepartment of Molecular Microbiology, Washington University Medical School, St. Louis, Missouri, USA

^bDepartment of Molecular and Cellular Biology, University of Guelph, Ontario, Canada

^cCenter for Bioinformatics, Faculty of Natural Sciences and Technology, Saarland University, Saarbruecken, Germany

^dDepartment of Biology, Faculty of Natural Sciences and Technology, Saarland University, Saarbruecken, Germany

ABSTRACT The fungal pathogen *Cryptococcus neoformans* is distinguished by a cell-wall-anchored polysaccharide capsule that is critical for virulence. Biogenesis of both cell wall and capsule relies on the secretory pathway. Protein secretion begins with polypeptide translocation across the endoplasmic reticulum (ER) membrane through a highly conserved channel formed by three proteins: Sec61, Sbh1, and Sss1. Sbh1, the most divergent, contains multiple phosphorylation sites, which may allow it to regulate entry into the secretory pathway in a species- and protein-specific manner. Absence of *SBH1* causes a cell-wall defect in both *Saccharomyces cerevisiae* and *C. neoformans*, although other phenotypes differ. Notably, proteomic analysis showed that when cryptococci are grown in conditions that mimic aspects of the mammalian host environment (tissue culture medium, 37°C, 5% CO₂), a set of secretory and transmembrane proteins is upregulated in wild-type, but not in $\Delta sbh1$ mutant cells. The Sbh1-dependent proteins show specific features of their ER targeting sequences that likely cause them to transit less efficiently into the secretory pathway. Many also act in cell-wall biogenesis, while several are known virulence factors. Consistent with these observations, the *C. neoformans* $\Delta sbh1$ mutant is avirulent in a mouse infection model. We conclude that, in the context of conditions encountered during infection, Sbh1 controls the entry of virulence factors into the secretory pathway of *C. neoformans*, and thereby regulates fungal pathogenicity.

IMPORTANCE *Cryptococcus neoformans* is a yeast that causes almost 200,000 deaths worldwide each year, mainly of immunocompromised individuals. The surface structures of this pathogen, a protective cell wall surrounded by a polysaccharide capsule, are made and maintained by proteins that are synthesized inside the cell and travel outwards through the secretory pathway. A protein called Sbh1 is part of the machinery that determines which polypeptides enter this export pathway. We found that when Sbh1 is absent, both *C. neoformans* and the model yeast *S. cerevisiae* show cell-wall defects. Lack of Sbh1 also changes the pattern of secretion of both transmembrane and soluble proteins, in a manner that depends on characteristics of their sequences. Notably, multiple proteins that are normally upregulated in conditions similar to those encountered during infection, including several needed for cryptococcal virulence, are no longer increased. Sbh1 thereby regulates the ability of this important pathogen to cause disease.

KEYWORDS *Cryptococcus neoformans*, *Saccharomyces cerevisiae*, Sbh1, cell wall, protein translocation, virulence

Cryptococcus neoformans is a haploid budding yeast that is ubiquitous in the environment, so that spores or desiccated cells are frequently inhaled (1, 2). In healthy individuals, these infectious particles are generally cleared or establish an asymptomatic latent infection (3). If, however, the host is or becomes severely immunocompromised, the pathogen may

Editor Alexander Idnurm, University of Melbourne

Copyright © 2023 Santiago-Tirado et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Karin Römisch, k.roemisch@mx.uni-saarland.de, or Tamara L. Doering, doering@wustl.edu.

*Present address: Felipe H. Santiago-Tirado, Department of Biological Sciences, University of Notre Dame, Notre Dame, Indiana, USA.

§Present address: Duy Nguyen, DKFZ Heidelberg, Germany.

The authors declare no conflict of interest.

This article is a direct contribution from Tamara L. Doering, a Fellow of the American Academy of Microbiology, who arranged for and secured reviews by Jeffrey Brodsky, University of Pittsburgh, and Marcio Rodrigues, Oswaldo Cruz Foundation (Fiocruz).

Received 9 December 2022

Accepted 23 December 2022

Published 7 February 2023




FIG 1 Sbh1 is required for cell wall integrity in *S. cerevisiae*. (A) Structure of the Sec61 channel of *S. cerevisiae* (modified from ref. 12; PDB 6ND1). The N-terminal half of Sec61 is shown in blue, C-terminal half in green, Sss1 in gray, and Sbh1 in magenta. The intrinsically unstructured N-terminal 38 amino acids of Sbh1 that were not visible in the cryo-EM structure were drawn in by hand. P, phosphorylation sites; dashed lines, position of the ER membrane; arrow, direction of protein transport. (B) Alignment of the poorly conserved N termini of Sbh1 proteins from *S. cerevisiae* (Sbh1, Sbh2) and the following pathogenic fungi: Cryn *Cryptococcus neoformans*, Sjap *Schizosaccharomyces japonicus*, Hcap *Histoplasma capsulatum*, Bder *Blastomyces dermatitidis*, Cpos *Coccidioides posadasii*, Afum *Aspergillus fumigatus*, Anig *Aspergillus niger*, Bfuc *Botryotinia fuckeliana*, Sbra *Sporothrix brasiliensis*. Except for Sbh2, only regions containing S or T residues with flanking prolines (pink) are shown. The conserved proline-flanked T5 in *S. cerevisiae* Sbh1 is indicated by an asterisk; other confirmed phosphorylation sites in *S. cerevisiae* Sbh1 are shown in gray; and the PKA site in *C. neoformans* Sbh1 is in blue. *Schizosaccharomyces cryophilus*, *Schizosaccharomyces octoporus*, *Candida albicans*, *Schizosaccharomyces pombe*, *Kluyveromyces lactis*, *Yarrowia lipolytica*, *Pichia pastoris*, and *Hansenula polymorpha* Sbh1 proteins were also screened, but do not contained potential proline-flanked phosphorylation sites. For full alignments of all Sbh1 protein sequences see Fig. S1. (C) Wild-type *SBH1 SBH2 S. cerevisiae* and the indicated mutants were grown on YPD plates at the indicated temperatures for 3 days. (D) The $\Delta sbh1\Delta sbh2$ *S. cerevisiae* strain was transformed with pRS415 without an insert (vector), wild-type *SBH1*, or the indicated phosphorylation site mutants and grown on SC without leucine containing 10 μ g/mL calcofluor white at 30°C. A $\Delta kre6$ mutant which is unable to grow on calcofluor white and its isogenic wild type were included as controls. Note that these are in a different strain background compared to the *SBH1* strain. (E) The indicated strains, including the untransformed $\Delta sbh1\Delta sbh2$ *S. cerevisiae* strain ($\Delta sbh1/2$), were grown to early exponential phase and cell-wall proteins were extracted by high pH/DTT, resolved by SDS-PAGE (size range shown is roughly 10–150 kD), and visualized by Coomassie staining.

grow and disseminate, both within and outside host cells (4). Dissemination to the central nervous system results in a devastating meningitis, which causes almost 200,000 deaths each year worldwide (5).

Like other fungi, *C. neoformans* is protected by a multilayer cell wall, which is composed of a meshwork of chitin, chitosan, alpha and beta glucans, and mannoproteins (6). This flexible and dynamic structure responds to and protects the cell from environmental stresses, while accommodating morphogenesis. An additional role of the cryptococcal wall is to anchor an elaborate polysaccharide capsule that surrounds the cell (7, 8). This structure, which is unique among fungal pathogens, is required for cryptococcal virulence (1). Capsule polysaccharides are also shed from the cell into the surrounding environment, where they act to perturb the host immune response (9, 10).

The biogenesis of both cell wall and capsule in *C. neoformans* relies on the secretory pathway, which begins with protein translocation across the ER membrane through the highly conserved Sec61 channel (11). An N-terminal hydrophobic signal peptide or a transmembrane domain targets proteins to this channel, which is formed by three proteins: Sec61, Sbh1, and Sss1 in yeast (Fig. 1A) and Sec61 α , Sec61 β , and Sec61 γ in mammals (11, 12). *SEC61* and *SSS1* are essential in *S. cerevisiae* (13, 14). The 10 transmembrane helices of Sec61 form a protein-conducting channel in the ER membrane, stabilized by Sss1 which

clamps around the Sec61 helix-bundle (Fig. 1A) (12, 15, 16). *SBH1* and its paralog *SBH2* are not essential, but deletion of both genes leads to temperature-sensitivity at 37°C in *S. cerevisiae* (17, 18). Sbh1/Sec61 β is a tail-anchored protein associated peripherally with the Sec61 channel (19) (Fig. 1A). Its conserved transmembrane domain is sufficient to rescue the double mutant growth defect at 37°C, while the role of its divergent cytosolic domain containing multiple phosphorylation sites is poorly understood (18, 20, 21). As is common for proteins with intrinsically unstructured domains, Sbh1/Sec61 β has multiple interactors: it binds to ribosomes and mediates association of the Sec61 channel with signal peptidase and signal recognition particle receptor (22–25). In mammalian cells, the Sbh1 homolog Sec61 β also interacts with the Stimulator of Interferon Genes (STING); this interaction is required for STING's ability to stimulate interferon expression, thus linking the Sec61 channel to the innate immune response (26). Sec61 with Sss1 can translocate proteins into the ER on its own, but the cytosolic domain of Sec61 β contacts secretory proteins prior to insertion in the cytosolic vestibule of the Sec61 channel and the presence of Sec61 β stimulates import (23, 27, 28). Recently, null mutations in human SEC61B were identified as a cause for polycystic liver disease (29). The mutations led to a complete biogenesis defect specific for polycystin-1, a large complex transmembrane protein with a weak secretory signal sequence (29). Collectively, these data have led to the view that Sbh1/Sec61 β mediates interaction of the Sec61 channel with other proteins in the ER membrane and that its cytosolic domain aids insertion of specific signal peptides or transmembrane domains into the Sec61 channel.

Phosphorylation is a frequent biological mechanism for regulation of protein activity or protein-protein interactions and a systematic analysis identified 63 kinases that play a role in the pathogenicity of *C. neoformans* (30, 31). One of these, Protein Kinase A (PKA), regulates the secretion of a subset of virulence factors (32). Regulation of protein secretion may be important during physiological transitions such as cell differentiation or development. Indeed Sec61 β —the only subunit of the Sec61 channel that has been found to be phosphorylated thus far—is important for polarization of epithelial cells, and for development of *Drosophila melanogaster* embryos (33–36). Both mammalian Sec61 β and yeast Sbh1 are phosphorylated at multiple sites in various combinations (21, 33; phosphosite databases). The phosphorylation sites all lie in the intrinsically unstructured cytosolic domain, and are not positionally conserved, with one exception: the proline-flanked site at T5 in *S. cerevisiae* is present in mammals and some birds, although not in lower vertebrates, invertebrates, or other nonpathogenic yeast species. This suggests that this phosphorylation site evolved at least twice by convergent evolution and thus likely has an important function (Fig. 1A; asterisk Fig. 1B) (21).

The 63 kinases required for virulence of *C. neoformans* include many proline-directed kinases (31, 37). When we aligned the N termini of Sbh1 proteins from nonpathogenic and pathogenic yeast, we noticed that—with the exception of *Candida albicans*—all pathogen Sbh1 proteins contain one or more potential proline-flanked phosphorylation site(s) in the N-terminal cytosolic domain, whereas Sbh1 proteins from nonpathogenic yeast species (*Schizosaccharomyces pombe*, *Kluyveromyces lactis*, *Yarrowia lipolytica*, *Pichia pastoris*, *Hansenula polymorpha*) and the *S. cerevisiae* homolog Sbh2 did not (Fig. 1B, pink; see Fig. S1 for full alignments). This led us to hypothesize that these phosphorylation sites in Sbh1 might be connected to virulence of the respective pathogenic species. In *C. neoformans* Sbh1, the proline-flanked phosphorylation sites are preceded by a potential PKA phosphorylation site (Fig. 1B, blue) and secretion of a subset of virulence factors in *C. neoformans* depends on PKA (32). We therefore decided to investigate the role of Sbh1 and its N-terminal phosphorylation sites in virulence of *C. neoformans*.

We found that, unlike in *S. cerevisiae*, mutations of the N-terminal Sbh1 phosphorylation sites (individually or in combination) do not affect *C. neoformans* growth in rich medium (YPD), and that even complete deletion of *SBH1* only inhibits growth when combined with high temperature and or cell-wall stressors. We also observed that in infection-like conditions the deletion mutant fails to properly regulate protein secretion. We have identified the Sbh1-dependent substrates and identified common characteristics of their ER-targeting sequences. Notably, the affected proteins include multiple virulence factors and the mutant is consequently avirulent in a mouse model of infection. Together, our results show that




FIG 2 Sbh1 is required for cell wall integrity at high temperature in *C. neoformans*. (A) Serial 10-fold dilutions of wild-type (KN99) and *sbh1*Δ cells were grown at the temperatures shown on YPD alone or containing 1 mg/mL calcofluor white (CFW) or 0.02% SDS (SDS). (B) Wild-type (KN99) or *sbh1*Δ mutant cells were transformed with vector alone (vec) or plasmids encoding HA-tagged Sbh1 (SBH1), or Sbh1 with mutations in the indicated phosphorylation sites and then diluted as in panel A and grown at the indicated conditions. (C) Cells of the strains indicated were treated with lysing enzymes (see Materials and Methods) for the times shown and then assessed for % lysis with two-way ANOVA with mixed-effects analysis. * indicates $P < 0.05$; # indicates $P = 0.25$. (D) The indicated strains were grown as above in the presence of 1 mM H_2O_2 or 1 mM $NaNO_2$ on YNB (pH 5) at 37°C. All experiments are representative of at least three biological replicate studies.

Sbh1 is critical for secretion of a subset of virulence factors and other proteins during *C. neoformans* infection, and is therefore essential for its pathogenicity.

RESULTS

Sbh1 is required for cell-wall integrity in *S. cerevisiae* and *C. neoformans*. We first looked to the best characterized system, *S. cerevisiae*. In this organism, both genes that encode the Sec61 channel beta subunit (*SBH1* and *SBH2*) are required for growth on YPD at 37°C (Fig. 1C) (17, 18, 21). *S. cerevisiae* lacking *SBH1* and *SBH2* are also resistant to calcofluor white (Fig. 1D), which suggests either reduced chitin in the cell-wall or altered cell-wall architecture. Further, we observed increased extractability of proteins from the cell wall by alkaline buffer containing dithiothreitol, which also suggests structural defects in the wall (Fig. 1E). While mutation of the conserved Sbh1 phosphorylation site T5 to A on its own has no effect on *S. cerevisiae* growth or cell-wall integrity (21), mutation of both proline-flanked phosphorylation sites to A phenocopied the absence of Sbh1 in *S. cerevisiae*. Mutation of both residues to phosphomimetic E had no effect (Fig. 1D and E).

The phenotypes we observed in *C. neoformans* cells lacking their sole *SBH1* gene were quite distinct from those of *S. cerevisiae*. Specifically, they showed only subtle temperature sensitivity at 37°C on YPD (Fig. 2A, left panel) and no sensitivity to osmotic or cell-wall stressors at 30°C (Fig. 2A). At 37°C, however, the mutant cells were hypersensitive to SDS and barely grew on calcofluor white (Fig. 2A, lower panels; Fig. 2B, top two rows); they were also more sensitive than wild-type cells to cell-wall digestion by lysing enzymes (Fig. 2C) and showed increased sensitivity to oxidative stress (1 mM H_2O_2 ; Fig. 2D, top two rows), which is encountered in the setting of infection. Overall, we conclude that Sbh1 is required for cell-wall integrity

in both the ascomycete *S. cerevisiae* and the basidiomycete *C. neoformans*, although specific manifestations of this requirement differ (see Discussion).

Sbh1 N-terminal phosphorylation sites do not affect cell-wall integrity in *C. neoformans*. As discussed above, we hypothesized that the N-terminal phosphorylation sites in cryptococcal Sbh1 were important for protein function. To test this, we first engineered a plasmid to express wild-type *SBH1* modified with a C-terminal HA tag. The tagged protein effectively complemented the sensitivity of *sbh1*Δ mutants to cell-wall stressors (CFW and SDS; Fig. 2B). We then modified this expression plasmid to encode Sbh1 that was altered to eliminate either the putative PKA site (Ser8 to Ala), two proline-flanked phosphorylation sites (Ser9 to Ala and Thr12 to Val), or all three sites. All of these polypeptides were expressed at similar levels, except for the version with a sole PKA site mutation, which was slightly less abundant (Fig. S2). We found that each of the mutated proteins fully complemented growth of *sbh1*Δ cells in the presence of CFW and also restored growth at 37°C in the presence of SDS (Fig. 2B). Interestingly, cells expressing mutant versions of Sbh1 grew slightly better on YPD and were more resistant to SDS than those expressing the WT protein (Fig. 2B). Each mutant protein also completely complemented the defective growth of the *sbh1*Δ mutant in the presence of oxidative stress (Fig. 2D). Together, these results suggest that, surprisingly, these phosphorylation sites are not required for Sbh1 function, as assessed by its role in cell-wall integrity.

Sbh1-dependent secretory and transmembrane proteins are induced in conditions that mimic the host environment. Sbh1 contacts signal peptides in the cytosolic vestibule of the Sec61 channel, so it can both guide secretory proteins into the channel and control their access to the secretory pathway. To address whether biogenesis of specific secretory or transmembrane proteins was compromised in the absence of *C. neoformans* Sbh1, we analyzed the proteomes of wild-type and *sbh1*Δ cells grown in either rich medium (YPD) at 30°C or in DMEM at 37°C and 5% CO₂. The latter, termed 37D5 below, was chosen to mimic key features of the complex host environment. We detected no significant differences between wild-type KN99 and *sbh1*Δ strains in YPD at 30°C (Fig. 3), suggesting no critical role for Sbh1 in those conditions. In 37D5, however, we discovered a set of proteins that was specifically induced in wild-type cells, but not in the *sbh1*Δ mutant (Fig. 3; Table S1). This set included 32 proteins without secretory pathway targeting sequences, 21 transmembrane (TM) proteins, and 68 proteins with signal peptides (SP) or uncleaved signal anchors (SA) (Table S1). When we analyzed their ER-targeting sequences, we found that the transmembrane domains were slightly more hydrophobic than average for transmembrane proteins in *C. neoformans*; many also had a strong positively charged patch in close proximity to one side of the TM domain (Fig. 4A; Table S2). Some of the transmembrane domains of the Sbh1-dependently induced proteins also contained a high number of prolines and/or glycines in addition to the polybasic stretch adjacent to the transmembrane domain (Table S2). The Sbh1-dependent signal peptides were more heterogeneous, but most displayed reduced C-region polarity (Fig. 4B; Table S2). In addition, they were suboptimal in various ways which therefore defied bioinformatic analysis: some had limited hydrophobicity, some contained high numbers of proline and glycine residues which interfere with alpha-helix formation, and many had a charge bias problem (either no charge on either side of the hydrophobic core, a positive charge C-terminal instead of N-terminal of the hydrophobic core, or multiple positive charges at the N-terminus of the mature part of the protein) (Table S2). Because signal peptides insert as alpha-helices into the hydrophobic lateral gate of the Sec61 channel with the positively charged N-terminus toward the cytosol, any of these features will reduce translocation efficiency into the ER. We suggest that Sbh1 is required for the biogenesis of specific secretory and transmembrane proteins under host-like conditions. Our model is that because the targeting sequences of these proteins are suboptimal for Sec61 channel insertion, they rely on the enhanced insertion efficiency provided by the presence of Sbh1 (Fig. 4C) (23).

Sbh1 is essential for virulence of *C. neoformans*. The proteome analysis presented above shows that Sbh1 regulates the secretion of proteins that are induced by host-like conditions. This process is likely important for pathogenesis, as supported by our observation that *sbh1*Δ cells have a reduced ability to survive within host macrophages (Fig. 5A and B). To directly test the role of Sbh1 in virulence, we used a well-established mouse model for




FIG 3 *C. neoformans* Sbh1 is required for the induction of secretory and transmembrane proteins in conditions that mimic the host. Proteomic analysis of wild-type and *sbh1*Δ *C. neoformans* grown in YPD at 30°C in room air or in DMEM at 37°C and 5% CO₂. Samples were analyzed in quadruplicate. Gene IDs of known virulence factors are indicated by asterisks.

cryptococcal infection. At 14 days after intranasal infection, we observed a striking reduction in both lung and brain fungal burden in *sbh1*Δ-infected animals compared to wild type (Fig. 5C and D). Consistent with this finding, mice infected with wild-type cryptococci succumbed to infection within 18 days, while mutant-infected animals remained healthy and grew normally for at least 10 weeks (Fig. 5E and F). We conclude that *SBH1* is essential for *C. neoformans* virulence.




FIG 4 Secretory and transmembrane proteins that require Sbh1 for induction contain specific ER-targeting sequences. (A) Characterization of Sbh1-dependent transmembrane domains. Top, schematic of a typical transmembrane domain (TMD) with features specific for Sbh1-dependent TMDs highlighted in red. Hydrophobicity (left) and charges within 10 amino acids of TMDs (right) were analyzed for TMDs of Sbh1 dependently induced proteins (candidates, red) compared to all transmembrane proteins in the *C. neoformans* proteome (black). (B) Characterization of Sbh1-dependent signal peptides. The schematic on top shows a typical signal peptide (SP) with features specific for Sbh1-dependent SPs highlighted in red. Black triangle: signal peptidase cleavage site. SPs of Sbh1-dependently induced secretory proteins (red) were analyzed for N-region charge, H-region hydrophobicity and length, and C-region polarity and compared to all signal peptides in the *C. neoformans* proteome (black). The only difference found was in C-region polarity, shown in the graph. (C) Model for Sbh1 function in *C. neoformans*. Under rich medium conditions, proteins with Sbh1-dependent ER targeting sequences (magenta) play only a minor role for vegetative growth (left), and most proteins imported into the ER have Sbh1-independent targeting sequences (shaded gray). During infection (right), however, Sbh1-dependent ER import (magenta) becomes essential.

DISCUSSION

In this work, we sought to identify factors regulating secretion of virulence-relevant proteins in the human pathogen *C. neoformans*. We specifically investigated the role of the nonessential ER protein translocation channel subunit Sbh1. Because the Sbh1 cytosolic domain contacts secretory signal sequences in the cytosolic vestibule of the channel prior to insertion into the channel, it can potentially aid or restrict protein import into the secretory pathway of specific proteins under specific growth conditions. This may be regulated by phosphorylation as Sbh1 in all species contains multiple phosphorylation sites which are only partially positionally conserved (Fig. 1A and B).




FIG 5 The *C. neoformans sbh1Δ* mutant is defective in growth in conditions that mimic the host or within host cells and is avirulent in mouse infection. (A) Fungi of the indicated strains were incubated in tissue culture medium (5% CO₂, 37°C) for 1, 18, or 30 h and then plated on YPD for enumeration of CFU (CFU; shown normalized to the 1 h CFU value). Shown are mean and SEM values from 4 independent experiments. (B) Cells were incubated for 1 h with differentiated THP-1 cells (MOI = 1), during which time fungi were engulfed by the host cells. Samples were then vigorously washed to remove free cryptococci and host cells were lysed to measure CFU either immediately or after 30 h of incubation. % change was calculated as ((Final CFU – Initial CFU)/Initial CFU) × 100. Shown are mean and SEM values from two independent experiments. For (A) and (B), * indicates *P* < 0.05 for *sbh1Δ* compared to KN99. (C and D) Total fungal burden at 14 days postinfection with the indicated strains, measured in lung (C) and brain (D). Each symbol represents CFU for one mouse; mean and standard deviation are also shown. Dotted line, original inoculum. (E) Mouse survival after intranasal infection with 5 × 10⁴ cells of the indicated strains. (F) Weight (mean and SD) of the mice from Panel E over time after infection. Any mouse that decreased to less than 80% of its initial weight (indicated by dashed lines) was humanely sacrificed and considered not to survive the infection.

In *S. cerevisiae*, absence of *SBH1* and its paralog *SBH2* causes temperature sensitivity, although the ER translocation defects at the restrictive temperature are modest for most substrates investigated (Fig. 1C) (18, 38). We hypothesized that this phenotype could relate to defects in the cell wall for several reasons: other translocation mutants exhibit cell-wall defects (39); wall synthesis relies on secretory proteins; and alterations in this process often yield a temperature-sensitive phenotype (40). We investigated the

effect of *sbh1* mutants on cell-wall integrity by assessing growth on calcofluor white (CFW; Fig. 1D) and extractability of cell-wall proteins by high pH/dithiothreitol treatment (Fig. 1E). CFW binds to chitin in the yeast cell wall and is toxic (41). *KRE6* encodes a glucosyl hydrolase required for beta-1,6-glucan synthesis; the deletion mutant is hypersensitive to CFW and is shown as a control. Cells lacking Sbh1 were slightly resistant to CFW (Fig. 1D), indicating a reduced amount of chitin in the cell wall, and displayed increased extraction of cell-wall proteins (Fig. 1E). *S. cerevisiae* cell-wall integrity also specifically required the two proline-flanked N-terminal phosphorylation sites at S3 and T5; mutation of these residues to nonphosphorylatable A phenocopied the cell-wall defects of the deletion mutant (Fig. 1D and E). The *sbh1* S3E/T5E mutant behaved like wild-type (Fig. 1D and E). Proline-flanked phosphorylation sites in Sbh1 of other yeast species occur primarily in human pathogens (Fig. 1B), which suggested to us that phosphorylation at these sites may play a regulatory role in virulence. The absence of *SBH1* in *C. neoformans* resulted only in a very modest temperature sensitivity at 37°C (Fig. 1C versus Fig. 2A); although at high temperature, *sbh1*Δ *C. neoformans* also displayed a cell-wall defect (Fig. 2A to C). In contrast to *S. cerevisiae*, however, mutation of the N-terminal proline-flanked phosphorylation sites of *C. neoformans* Sbh1 individually or in combination did not phenocopy the cell-wall defect (Fig. 2B). As protein kinase A has been shown to be critical for *C. neoformans* virulence, and *C. neoformans* Sbh1 also contains a PKA site (Fig. 1B, blue), we also mutated this site alone or in combination with the proline-flanked sites, but again saw no effect on cell-wall integrity (Fig. 2B) (32). Surprisingly, the altered phosphorylation sites slightly enhanced cell growth, especially notable on SDS (Fig. 2B). Taken together, our data suggest that Sbh1 is important for efficient secretion of one or more enzymes or precursors for the cell wall in both *S. cerevisiae* and in *C. neoformans*, but that the role of the N-terminal proline-flanked phosphorylation sites in this process differs in the two organisms. Alternatively, the differential effect of the phosphorylation site mutations may reflect their distinct cell-wall compositions: whereas the *S. cerevisiae* cell wall is formed by β-1,3-glucan, β-1,6-glucan, chitin, and mannoproteins, in the *C. neoformans* cell wall, most chitin is deacylated to chitosan, and α-1,3-glucan is a major cell-wall component (10, 42).

We next asked which proteins were affected by absence of *SBH1* in *C. neoformans*. Using a proteomic approach, we identified a set of secretory and transmembrane proteins that were induced under 37D5 conditions in the wild-type strain, but not in the *sbh1*Δ mutant (Fig. 3; Tables S1 and S2). Many of the membrane proteins in this set were characterized by a strong polybasic patch adjacent to the first transmembrane domain (e.g., CNAG_1854, CNAG_5502, CNAG_6416), while the signal peptides of the secretory proteins were more heterogeneous, and had less polar C-regions than average (Table S2; Fig. 4A and B). This is reminiscent of the ER targeting signals identified by Ziska and colleagues in mammalian cells whose ER import was dependent on the Sec61 channel-interacting protein Sec63 and the ER-luminal Hsp70 chaperone BiP (43). In *S. cerevisiae* the Sec63 complex composed of Sec63, Sec62, Sec71, and Sec72 promotes posttranslational protein import into the ER through the Sec61 channel by stabilizing the lateral gate in a partially open conformation (12, 44). It also plays a role in membrane protein topogenesis in both yeast and mammalian cells (45, 46). Sec63 is a polytopic membrane protein with an ER-luminal J-domain that activates ATP-hydrolysis of the ER-luminal chaperone BiP (Kar2 in yeast) (47). Kar2 contributes to protein import into the ER either by directly binding to the incoming protein or by promoting its folding (48, 49). As proteins dependent on Sbh1 in *C. neoformans* and proteins dependent on Sec63/BiP in mammalian cells have similar characteristics, our data suggest that Sbh1 and the Sec63 complex have overlapping functions in promoting ER import of specific proteins.

Signal peptides generally insert into the Sec61 channel with their N-terminus toward the cytosol and their N-terminal positive charge (Fig. 4B; “+”) contributes to this orientation (50). Many of the Sbh1-dependent signal peptides we identified in *C. neoformans* have no charge bias toward the N-terminus; in some, the charge bias is even reversed (Table S2). Sbh1 may help orient such signal peptides in the cytosolic vestibule of the Sec61 channel with the help of its cytosolic domain. The polar C-region of signal peptides contains the signal peptidase cleavage site (50). Most of the Sbh1-dependent signal peptides we identified have a very short C-region (Table S2), which may limit signal peptidase access unless the signal peptide

TABLE 1 Sbh1-dependent proteins involved in cryptococcal virulence^a

| Gene ID | Gene name | Protein function | SP or TM | WT/ <i>sbh1</i> Δ proteomics (log ₂ -fold) | Virulence citations | 24 h/0 h RNA-seq |
|------------|--------------|--|----------|---|---------------------|------------------|
| CNAG_00150 | | Serine endopeptidase, cell wall | SP1-23 | 5.24 | 53 | 1.68 |
| CNAG_01653 | <i>CIG1</i> | Cytokine-inducing glycoprotein | SP1-19 | 4.00 | 68 | 385.34 |
| CNAG_01172 | <i>PBX1</i> | Beta-helix containing protein | SP1-18 | 3.68 | 55, 53, 56 | 0.78 |
| CNAG_04735 | <i>MPR1</i> | Extracellular metalloproteinase (Mep1) | SP1-19 | 3.44 | 53, 59 | 122.79 |
| CNAG_00897 | <i>SKN1</i> | Glucosidase | TM134-58 | 3.04 | 57 | 3.43 |
| CNAG_02030 | | Glyoxal oxidase | SP1-21 | 2.79 | 53 | 0.97 |
| CNAG_00600 | <i>CAP60</i> | Capsule-associated protein | SP1-33 | 2.65 | 58 | 1.97 |
| CNAG_01331 | | | SP1-18 | 2.62 | 53 | 13.45 |

^aThese proteins meet the following criteria: their expression in 37D5 was at least 2-fold higher in WT compared to *sbh1*Δ cells; they contain an SP or TM domain; and they are required for normal *C. neoformans* virulence in mouse infections. Note that this is likely a subset of such proteins, as most cryptococcal proteins have not been tested for roles in virulence; other candidates that meet the first two criteria are listed in Table S1. Proteomics results are shown as the ratio of expression in WT versus *sbh1*Δ cells in 37D5. RNA-seq results are shown as the ratio of gene expression at 24 h versus 0 h in 37D5 (61). Expression of a subset of these genes (CNAG_04735, CNAG_02030, CNAG_00897) was measured by qPCR; these studies confirmed that it is not affected by loss of *SBH1* (Fig. S3).

or signal peptidase are positioned accurately by Sbh1. In order to efficiently insert into the Sec61 channel, the signal peptides of secretory proteins require an alpha-helical hydrophobic region (Fig. 4B and H, blue) (50). Many of the Sbh1-dependent signal peptides of *C. neoformans* contain multiple helix-breaking P or G residues within their hydrophobic helical core regions, and some have very short or very long hydrophobic H-regions (Table S2). These features would make it difficult for signal peptides to stably insert into the Sec61 channel lateral gate. Overexpression of an N-terminally truncated Sbh1 lacking most of its cytosolic domain complements the temperature-sensitivity and at least some of the translocation defect in *Δsbh1 Δsbh2 S. cerevisiae* (18). Together with our data, this suggests that the Sbh1 transmembrane domain stabilizes a conformation of the Sec61 lateral gate that favors insertion of the helical hydrophobic core of signal peptides, even if they are suboptimal.

That Sbh1-dependent secretion is critical for virulence is illustrated by both the reduced intracellular fitness of *sbh1*Δ *C. neoformans* in macrophages and the effects in mice. In contrast to infection with KN99 wild-type *C. neoformans*, which resulted in a high burden in the lungs and death of all mice by day 18 postinfection, all mice infected with *sbh1*Δ *C. neoformans* survived for 70 days and even gained weight (Fig. 5), showing that the mutant strain was avirulent.

We hypothesize that the striking defect in virulence of the *sbh1* mutant reflects reduced secretion of multiple groups of proteins. For example, these could include proteins required for fungal growth in the host environment (e.g., metabolic genes or proteins involved in cell-wall synthesis), needed for virulence factor production, or directly involved in host damage. These categories may also interact; for example, mutations that alter cell-wall glycan synthesis also result in aberrant formation of the capsule (7, 8, 51, 52). To explore this idea, we reviewed the polypeptides whose secretion is most impaired in the absence of Sbh1 (Table S1), focusing on proteins with signal or transmembrane sequences that had been previously implicated in cryptococcal virulence. We identified eight such proteins, which had exhibited infectivity defects in a large-scale screen (53) and/or been individually studied and shown to contribute to virulence (Table 1). One of them, *Cig1*, has also been detected in the blood of mice infected with *C. neoformans*, leading to the suggestion that it could serve as a biomarker of infection (32). Notably, all of the corresponding genes are expressed in human CSF during cryptococcal meningitis, most at higher levels than in rich medium (Table 1 and [54]). Two proteins in this group (*Pbx1* [55, 56] and *Skn1* [57]) act in cell-wall synthesis, so their reduced secretion could contribute to the wall defects and sensitivity to host-induced stress that we observed in *sbh1* mutant cells. Another is required for capsule display (*Cap60* [58]), although its biochemical function has not been characterized, and two others are proteases (CNAG_00150 and *Mpr1* [59]). Many fungal proteases damage host tissues during infection (60) and *Mpr1* further acts in cryptococcal crossing of the blood-brain barrier (59), which is required for cryptococcal dissemination to the brain and subsequent lethality. Supporting a role in infection for this group of proteins, six of the corresponding genes are expressed more highly in 37D5 than in rich medium (one other is modestly decreased); this pattern is particularly striking for *CIG1* and *MPR1*, whose expression levels increase over 100-fold (Table 1) (61).

Notably, the evidence that these proteins are important for cryptococcal virulence derives from studies of individual gene deletion strains, which completely lack one of these target proteins. The situation in the *sbh1* mutant is different, in that many proteins are secreted less efficiently, but none (besides Sbh1) are absent. Nonetheless, the impaired secretion of multiple proteins with key roles in virulence, together with others in the functional groups above, provides abundant reason for the inability of *sbh1* to cause disease in a mammalian host.

We conclude that *C. neoformans* Sbh1 is important for biogenesis of virulence factors whose specific ER-targeting sequences make these proteins dependent on Sbh1 for ER import (Fig. 4C). Recently, Sec61beta has also been shown to be critical for infection by the important fungal crop pathogen *Magnaporthe oryzae*, suggesting a more general role for Sbh1/Sec61beta in fungal infection (62). Because the intrinsically unstructured N-terminal part of the Sbh1 cytosolic domain is not conserved between yeast and mammals, our work may suggest a new specific target for developing drugs against this important human pathogen.

MATERIALS AND METHODS

***S. cerevisiae* methods.** *S. cerevisiae* strains are detailed in Text S1 and were grown on either yeast peptone dextrose (YPD) or synthetic complete medium without leucine, with or without 10 μ g/mL calcofluor white (Sigma), at 30°C or 37°C, for 3 days. Cell-wall proteins were extracted from cells in early exponential phase in 100 mM Tris-HCl, pH 9.4, 10 mM DTT at 37°C, and TCA-precipitated for SDS-PAGE analysis as detailed in Text S1.

***C. neoformans* strains, growth conditions, and plasmids.** All strains were in the *C. neoformans* serotype A strain KN99 α background, as detailed in Text S1, and were grown at 30°C on YPD with 100 μ g/mL of nourseothricin or 100 μ g/mL G418 as appropriate. After plasmids expressing the native *SBH1* gene or a version with 5 HA epitopes at the C-terminus were shown to complement phenotypic defects of the *sbh1* mutant, the tagged version was mutagenized to modify N-terminal motifs and electroporated into KN99 α and the *sbh1* Δ strains as described in (63) and detailed in Text S1. Because of the heterogeneity in copy number associated with plasmids in *C. neoformans*, at least three independent colonies were picked and tested for each construct; all exhibited the same behaviors.

Phenotypic assays and mass spectrometry. Serial dilution plate assays and sensitivity to hypotonic lysis after exposure to lysing enzyme were assessed as detailed in Text S1 and (64). Total cellular proteome analysis was performed as in (65, 66) and Text S1. Results were compared to the *C. neoformans* reference proteome and deposited in the PRIDE partner repository for the ProteomeXchange Consortium with the data set identifier: [PXD013894](https://www.ebi.ac.uk/pride/projects/PXD013894). A Student's *t* test was performed to identify proteins with significant differential expression (*P*-value < 0.05) ($S_0 = 1$) between samples employing a 5% permutation-based FDR filter.

Infection assays. Fungal survival after engulfment by THP-1 (ATCC #TIB-202) cells *in vitro* was assessed as in (67), with minor modifications detailed in Text S1. For *in vivo* studies, 4- to 6-week-old female C57BL/6 mice were intranasally inoculated with 5×10^4 cryptococcal cells and either followed by weight for survival studies or sacrificed at set times for organ burden measurement, as detailed in Text S1. All animal studies were reviewed and approved by the Animal Studies Committee of Washington University School of Medicine and conducted according to the National Institutes of Health guidelines for housing and care of laboratory animals.

Data availability. Raw data files are available in the PRIDE (Proteomics Identification Database); Project ID: [PXD013894](https://www.ebi.ac.uk/pride/projects/PXD013894).

SUPPLEMENTAL MATERIAL

Supplemental material is available online only.

TEXT S1, PDF file, 0.1 MB.

FIG S1, PDF file, 0.3 MB.

FIG S2, PDF file, 1 MB.

FIG S3, PDF file, 0.1 MB.

TABLE S1, XLSX file, 0.02 MB.

TABLE S2, DOCX file, 0.03 MB.

ACKNOWLEDGMENTS

We thank William Allen, Bristol University, for help with Fig. 1A; Christina Soromani, Andrea Ludes, and Nicole Spitzlberger (all former students of the Römisch lab) for data shown in Fig. 1C and D; Alyssa R. Brunsmann, Daphne Ko, and Liza Loza for qPCR studies; and Jussi Jääntti, VTT Helsinki, for *S. cerevisiae* strains and plasmids.

K.R. was funded by core funding of the Saarland University. Work in the Doering lab was supported by NIH grants AI140979 and AI135012 to T.L.D. F.H.S.-T. also received support from NIH T32 grant AI007172, a Postdoctoral Enrichment Program Award from the Burroughs Wellcome Fund, and NIH R21 grant AI171742 from NIAID. J.G.-M. is supported by the Canadian Foundation for Innovation (JELF #38798) and the Canadian Institutes for Health

Research (Project Grant). V.H. was supported by Deutsche Forschungsgemeinschaft grant He3875/15-1. The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

REFERENCES

- Kwon-Chung K, Fraser JA, Doering TL, Wang Z, Janbon G, Idnurm A, Bahn YS. 2014. *Cryptococcus neoformans* and *Cryptococcus gattii*, the etiologic agents of cryptococcosis. *Cold Spring Harb Perspect Med* 4:a019760. <https://doi.org/10.1101/cshperspect.a019760>.
- Srikanta D, Santiago-Tirado FH, Doering TL. 2014. *Cryptococcus neoformans*: historical curiosity to modern pathogen. *Yeast* 31:47–60. <https://doi.org/10.1002/yea.2997>.
- Montoya MC, Magwene PM, Perfect JR. 2021. Associations between cryptococcus genotypes, phenotypes, and clinical parameters of human disease: a review. *J Fungi (Basel)* 7:260. <https://doi.org/10.3390/jof7040260>.
- Gaylord EA, Choy HL, Doering TL. 2020. Dangerous liaisons: interactions of *Cryptococcus neoformans* with host phagocytes. *Pathogens* 9:891–905. <https://doi.org/10.3390/pathogens9110891>.
- Rajasingham R, Smith RM, Park BJ, Jarvis JN, Govender NP, Chiller TM, Denning DW, Loyse A, Boulware DR. 2017. Global burden of disease of HIV-associated cryptococcal meningitis: an updated analysis. *Lancet Infectious Dis* 17:873–881. [https://doi.org/10.1016/S1473-3099\(17\)30243-8](https://doi.org/10.1016/S1473-3099(17)30243-8).
- Loza L, Doering TL. 2021. Glycans of the pathogenic yeast *Cryptococcus neoformans* and related opportunities for therapeutic advances, p 479–506. *In* Barchi J. (ed) *Comprehensive Glycoscience*, 2nd ed, Elsevier, Amsterdam, the Netherlands.
- Reese AJ, Doering TL. 2003. Cell wall alpha-1,3-glucan is required to anchor the *Cryptococcus neoformans* capsule. *Mol Microbiol* 50:1401–1409. <https://doi.org/10.1046/j.1365-2958.2003.03780.x>.
- Reese AJ, Yoneda A, Bregger JA, Beauvais A, Liu H, Griffith CL, Bose I, Kim MJ, Skau C, Yang S, Sefko JA, Osumi M, Latge JP, Mylonakis E, Doering TL. 2007. Loss of cell wall alpha(1–3) glucan affects *Cryptococcus neoformans* from ultrastructure to virulence. *Mol Microbiol* 63:1385–1398. <https://doi.org/10.1111/j.1365-2958.2006.05551.x>.
- Agustinho DP, Miller LC, Li LX, Doering TL. 2018. Peeling the onion: the outer layers of *Cryptococcus neoformans*. *Mem Inst Oswaldo Cruz* 113:e180040. <https://doi.org/10.1590/0074-02760180040>.
- Wang ZA, Li LX, Doering TL. 2018. Unravelling synthesis of the cryptococcal cell wall and capsule. *Glycobiology* 28:719–730. <https://doi.org/10.1093/glycob/cwy030>.
- Voorhees R, Hegde RS. 2016. Toward a structural understanding of cotranslational protein translocation. *Curr Opin Cell Biol* 41:91–99. <https://doi.org/10.1016/j.cob.2016.04.009>.
- Wu X, Cabanos C, Rapoport TA. 2019. Structure of the post-translational protein translocation machinery of the ER membrane. *Nature* 566:136–139. <https://doi.org/10.1038/s41586-018-0856-x>.
- Stirling CJ, Rothblatt J, Hosobuchi M, Deshaies R, Schekman R. 1992. Protein translocation mutants defective in the insertion of integral membrane proteins into the endoplasmic reticulum. *Mol Biol Cell* 3:129–142. <https://doi.org/10.1091/mbc.3.2.129>.
- Esnault Y, Blondel MO, Deshaies RJ, Schekman R, Képès F. 1993. The yeast SSS1 gene is essential for secretory protein translocation and encodes a conserved protein of the endoplasmic reticulum. *EMBO J* 12:4083–4093. <https://doi.org/10.1002/j.1460-2075.1993.tb06092.x>.
- Esnault Y, Feldheim D, Blondel MO, Schekman R, Képès F. 1994. SSS1 encodes a stabilizing component of the Sec61 subcomplex of the yeast protein translocation apparatus. *J Biol Chem* 269:27478–27485. [https://doi.org/10.1016/S0021-9258\(18\)47010-X](https://doi.org/10.1016/S0021-9258(18)47010-X).
- Falcone D, Henderson MP, Nieuwland H, Coughlan CM, Brodsky JL, Andrews DW. 2011. Stability and function of the Sec61 translocation complex depends on the Sss1p tail-anchor sequence. *Biochem J* 436:291–303. <https://doi.org/10.1042/BJ20101865>.
- Toikkanen J, Gatti E, Takei K, Saloheimo M, Olkkonen VM, Söderlund H, de Camilli P, Keränen S. 1996. Yeast protein translocation complex: isolation of two genes *SEB1* and *SEB2* encoding proteins homologous to the Sec61 β subunit. *Yeast* 12:425–438. [https://doi.org/10.1002/\(SICI\)1097-0061\(199604\)12:5<425::AID-YEA924>3.0.CO;2-B](https://doi.org/10.1002/(SICI)1097-0061(199604)12:5<425::AID-YEA924>3.0.CO;2-B).
- Feng D, Zhao X, Soromani C, Toikkanen J, Römisch K, Vembar SS, Brodsky JL, Keränen S, Jäntti J. 2007. The transmembrane domain is sufficient for Sbh1p function, its association with the Sec61 complex, and interaction with Rtn1p. *J Biol Chem* 282:30618–30628. <https://doi.org/10.1074/jbc.M701840200>.
- Leznicki P, Warwicker J, High S. 2011. A biochemical analysis of the constraints of tail-anchored protein biogenesis. *Biochem J* 436:719–727. <https://doi.org/10.1042/BJ20101737>.
- Leroux A, Rokeach LA. 2008. Interspecies complementation of the translocon beta subunit requires only its transmembrane domain. *PLoS One* 3:e3880. <https://doi.org/10.1371/journal.pone.0003880>.
- Soromani C, Zeng N, Hollemeyer K, Heinze E, Klein M-C, Tretter T. 2012. N-acetylation and phosphorylation of Sec complex subunits in the ER membrane. *BMC Cell Biol* 13:34. <https://doi.org/10.1186/1471-2121-13-34>.
- Levy R, Wiedmann M, Kreibich G. 2001. *In vitro* binding of ribosomes to the β subunit of the Sec61p protein translocations complex. *J Biol Chem* 276:2340–2346. <https://doi.org/10.1074/jbc.M004867200>.
- Kalies K-U, Rapoport TA, Hartmann E. 1998. The β subunit of the Sec61 complex facilitates cotranslational protein transport and interacts with signal peptidase during translocation. *J Cell Biol* 141:887–894. <https://doi.org/10.1083/jcb.141.4.887>.
- Antonin W, Meyer HA, Hartmann E. 2000. Interactions between Spc2 and other components of the endoplasmic reticulum translocation sites of the yeast *Saccharomyces cerevisiae*. *J Biol Chem* 275:34068–34072. <https://doi.org/10.1074/jbc.M006126200>.
- Jiang Y, Cheng Z, Mandon EC, Gilmore R. 2008. An interaction between the SRP receptor and the translocon is critical during cotranslational protein translocation. *J Cell Biol* 180:1149–1161. <https://doi.org/10.1083/jcb.200707196>.
- Ishikawa H, Barber GN. 2008. STING is an endoplasmic reticulum adaptor that facilitates innate immune signalling. *Nature* 455:674–678. <https://doi.org/10.1038/nature07317>.
- Laird V, High S. 1997. Discrete cross-linking products identified during membrane protein biosynthesis. *J Biol Chem* 272:1983–1989. <https://doi.org/10.1074/jbc.272.3.1983>.
- Besemer J, Harant H, Wang S, Oberhauser B, Marquardt K, Foster SA, Schreiner EP, de Vries JE, Dascher-Nadel C, Lindley IJD. 2005. Selective inhibition of cotranslational translocation of vascular cell adhesion molecule 1. *Nature* 436:290–293. <https://doi.org/10.1038/nature03670>.
- Besse W, Dong K, Choi J, Punia S, Fedeles SV, Choi M, Gallagher AR, Huang EB, Gulati A, Knight J, Mane S, Tahvanainen E, Tahvanainen P, Sanna-Cherchi S, Lifton RP, Watnick T, Pei YP, Torres VE, Somlo S. 2017. Isolated polycystic liver disease genes define effectors of polycystin-1 function. *J Clin Invest* 127:1772–1785. <https://doi.org/10.1172/JCI90129>.
- Ubersax JA, Ferrell JE, Jr. 2007. Mechanisms of specificity in protein phosphorylation. *Nat Rev Mol Cell Biol* 8:530–541. <https://doi.org/10.1038/nrm2203>.
- Lee K-T, So Y-S, Yang D-H, Jung K-W, Choi J, Lee D-G, Kwon H, Jang J, Wang LL, Cha S, Meyers GL, Jeong E, Jin J-H, Lee Y, Hong J, Bang S, Ji J-H, Park G, Byun H-J, Park SW, Park Y-M, Adedoyin G, Kim T, Averette AF, Choi J-S, Heitman J, Cheong E, Lee Y-H, Bahn Y-S. 2016. Systematic functional analysis of kinases in the fungal pathogen *Cryptococcus neoformans*. *Nat Commun* 7:12766. <https://doi.org/10.1038/ncomms12766>.
- Geddes JMH, Croll D, Caza M, Stoynov N, Foster LJ, Kronstad JW. 2015. Secretome profiling of *Cryptococcus neoformans* reveals regulation of a subset of virulence-associated proteins and potential biomarkers by protein kinase A. *BMC Microbiol* 15:206. <https://doi.org/10.1186/s12866-015-0532-3>.
- Gruss OJ, Feick P, Frank R, Dobberstein B. 1999. Phosphorylation of components of the ER translocation site. *Eur J Biochem* 260:785–793. <https://doi.org/10.1046/j.1432-1327.1999.00215.x>.
- Choi SY, Fogelgren B, Zuo Z, Juang L, McKenna S, Lingappa VR, Lipschutz JH. 2012. Exocyst Sec10 is involved in basolateral protein translation and translocation in the endoplasmic reticulum. *Neuron Exp Nephrol* 120:e133–e139.
- Washiyama M, Koizumi N, Fujii M, Watanabe Y. 2014. Sec61 β regulates barrier functions of tight junctions through expression of claudin-4 in MDCK cells. *Biol Pharm Bull* 37:979–986. <https://doi.org/10.1248/bpb.13-00936>.
- Valcarcel R, Weber U, Jackson DB, Benes V, Ansorge W, Bohmann D, Mlodzik M. 1999. Sec61beta, a subunit of the protein translocation channel, is required during *Drosophila* development. *J Cell Sci* 112:4389–4396. <https://doi.org/10.1242/jcs.112.23.4389>.
- Kannan N, Neuwald AF. 2004. Evolutionary constraints associated with functional specificity of the CMGC protein kinases MAPK, CDK, GSK, SRPK,

- DYRK, and CK2alpha. *Protein Sci* 13:2059–2077. <https://doi.org/10.1110/ps.04637904>.
38. Finke K, Plath K, Panzner S, Prehn S, Rapoport TA, Hartmann E, Sommer T. 1996. A second trimeric complex containing homologs of the Sec61p complex functions in protein transport across the ER membrane of *S. cerevisiae*. *EMBO J* 15:1482–1494. <https://doi.org/10.1002/j.1460-2075.1996.tb00492.x>.
 39. Barbieri G, Simon J, Lupusella CR, Pereira F, Elia F, Meyer H, Schuldiner M, Hanes SD, Nguyen D, Helms V, Römisch K. 2023. Sbh1/Sec61 β promotes ER translocation of proteins with suboptimal targeting sequences and is fine-tuned by phosphorylation. *Biorxiv* <https://doi.org/10.1101/2022.05.18.492448>.
 40. Wright CM, Fewell SW, Sullivan ML, Pipas JM, Watkins SC, Brodsky JL. 2007. The Hsp40 molecular chaperone Ydj1p, along with the protein kinase C pathway, affects cell-wall integrity in the yeast *Saccharomyces cerevisiae*. *Genetics* 175:1649–1664. <https://doi.org/10.1534/genetics.106.066274>.
 41. Roncero C, Valdivieso MH, Ribas JC, Durán A. 1988. Isolation and characterization of *Saccharomyces cerevisiae* mutants resistant to Calcofluor white. *J Bacteriol* 170:1950–1954. <https://doi.org/10.1128/jb.170.4.1950-1954.1988>.
 42. Lesage G, Bussey H. 2006. Cell wall assembly in *Saccharomyces cerevisiae*. *Microbiol Mol Biol Rev* 70:317–343. <https://doi.org/10.1128/MMBR.00038-05>.
 43. Ziska A, Tatzelt J, Dudek J, Paton AW, Paton JC, Zimmermann R, Ha β denteufel S. 2019. The signal peptide plus a cluster of positive charges in prion protein dictate chaperone-mediated Sec61-channel gating. *Biol Open* 8:bio040691.
 44. Itskanov S, Park E. 2019. Structure of the posttranslational Sec protein translocation channel complex from yeast. *Science* 363:84–87. <https://doi.org/10.1126/science.aav6740>.
 45. Mades A, Gotthardt K, Awe K, Stieler J, Döring T, Füsler S, Prange R. 2012. Role of human Sec63 in modulating the steady-state levels of multi-spanning membrane proteins. *PLoS One* 7:e49243. <https://doi.org/10.1371/journal.pone.0049243>.
 46. Jung SJ, Kim JE, Reithinger JH, Kim H. 2014. The Sec62–Sec63 translocon facilitates translocation of the C-terminus of membrane proteins. *J Cell Sci* 127:4270–4278. <https://doi.org/10.1242/jcs.153650>.
 47. Vembar SS, Jonikas MC, Hendershot LM, Weissman JS, Brodsky JL. 2010. J domain co-chaperone specificity defines the role of BiP during protein translocation. *J Biol Chem* 285:22484–22494. <https://doi.org/10.1074/jbc.M110.102186>.
 48. Matlack KE, Misselwitz B, Plath K, Rapoport TA. 1999. BiP acts as a molecular ratchet during posttranslational transport of prepro-alpha factor across the ER membrane. *Cell* 97:553–564. [https://doi.org/10.1016/s0092-8674\(00\)80767-9](https://doi.org/10.1016/s0092-8674(00)80767-9).
 49. Allen WJ, Collinson I, Römisch K. 2019. Post-translational protein transport by the Sec complex. *Trends Biochem Sci* S0968-0004:30056–30058.
 50. Hegde RS, Bernstein HD. 2006. The surprising complexity of signal sequences. *Trends Biochem Sci* 31:563–571. <https://doi.org/10.1016/j.tibs.2006.08.004>.
 51. Banks IR, Specht CA, Donlin MJ, Gerik KJ, Levitz SM, Lodge JK. 2005. A chitin synthase and its regulator protein are critical for chitosan production and growth of the fungal pathogen *Cryptococcus neoformans*. *Eukaryot Cell* 4:1902–1912. <https://doi.org/10.1128/EC.4.11.1902-1912.2005>.
 52. Baker LG, Specht CA, Donlin MJ, Lodge JK. 2007. Chitosan, the deacetylated form of chitin, is necessary for cell wall integrity in *Cryptococcus neoformans*. *Eukaryot Cell* 6:855–867. <https://doi.org/10.1128/EC.00399-06>.
 53. Liu OW, Chun CD, Chow ED, Chen C, Madhani HD, Noble SM. 2008. Systematic genetic analysis of virulence in the fungal pathogen *Cryptococcus neoformans*. *Cell* 135:174–188. <https://doi.org/10.1016/j.cell.2008.07.046>.
 54. Chen Y, Toffaletti DL, Tenor JL, Litvintseva AP, Fang C, Mitchell TG, McDonald TR, Nielsen K, Boulware DR, Bicanic T, Perfect JR. 2014. The *Cryptococcus neoformans* transcriptome at the site of human meningitis. *mBio* 5:e01087-13–e01013. <https://doi.org/10.1128/mBio.01087-13>.
 55. Liu OW, Kelly MJ, Chow ED, Madhani HD. 2007. Parallel beta-helix proteins required for accurate capsule polysaccharide synthesis and virulence in the yeast *Cryptococcus neoformans*. *Eukaryot Cell* 6:630–640. <https://doi.org/10.1128/EC.00398-06>.
 56. Kumar P, Heiss C, Santiago-Tirado FH, Black I, Azadi P, Doering TL. 2014. Pbx proteins in *Cryptococcus neoformans* cell wall remodeling and capsule assembly. *Eukaryot Cell* 13:560–571. <https://doi.org/10.1128/EC.00290-13>.
 57. Gilbert NM, Donlin MJ, Gerik KJ, Specht CA, Djordjevic JT, Wilson CF, Sorrell TC, Lodge JK. 2010. *KRE* genes are required for beta-1,6-glucan synthesis, maintenance of capsule architecture and cell wall protein anchoring in *Cryptococcus neoformans*. *Mol Microbiol* 76:517–534. <https://doi.org/10.1111/j.1365-2958.2010.07119.x>.
 58. Chang YC, Kwon-Chung KJ. 1998. Isolation of the third capsule-associated gene, *CAP60*, required for virulence in *Cryptococcus neoformans*. *Infect Immun* 66:2230–2236. <https://doi.org/10.1128/IAI.66.5.2230-2236.1998>.
 59. Vu K, Tham R, Uhrig JP, Thompson GR, 3rd, Na Pombejra S, Jamklang M, Bautos JM, Gelli A. 2014. Invasion of the central nervous system by *Cryptococcus neoformans* requires a secreted fungal metalloprotease. *mBio* 5:e01101-14–e01114. <https://doi.org/10.1128/mBio.01101-14>.
 60. Monod M, Capoccia S, Lechenne B, Zaugg C, Holdom M, Jousson O. 2002. Secreted proteases from pathogenic fungi. *Int J Med Microbiol* 292:405–419. <https://doi.org/10.1078/1438-4221-00223>.
 61. Maier EJ, Haynes BC, Gish SR, Wang ZA, Skowrya ML, Marulli AL, Doering TL, Brent MR. 2015. Model-driven mapping of transcriptional networks reveals the circuitry and dynamics of virulence regulation. *Genome Res* 25:690–700. <https://doi.org/10.1101/gr.184101.114>.
 62. Wei YY, Liang S, Zhang YR, Lu JP, Lin FC, Liu XH. 2020. MoSec61 β , the beta subunit of Sec61, is involved in fungal development and pathogenicity, plant immunity, and ER-phagy in *Magnaporthe oryzae*. *Virulence* 11:1685–1700. <https://doi.org/10.1080/21505594.2020.1848983>.
 63. Skowrya ML, Doering TL. 2012. RNA interference in *Cryptococcus neoformans*. *Methods Mol Biol* 845:165–186. https://doi.org/10.1007/978-1-61779-539-8_11.
 64. Gerik KJ, Donlin MJ, Soto CE, Banks AM, Banks IR, Maligie MA, Selitrennikoff CP, Lodge JK. 2005. Cell wall integrity is dependent on the *PKC1* signal transduction pathway in *Cryptococcus neoformans*. *Mol Microbiol* 58:393–408. <https://doi.org/10.1111/j.1365-2958.2005.04843.x>.
 65. Geddes JM, Caza M, Croll D, Stoyanov N, Foster LJ, Kronstad JW. 2016. Analysis of the protein kinase A-regulated proteome of *C. neoformans* identifies a role for the ubiquitin-proteasome pathway in capsule formation. *mBio* 7:e01862-15–e01815. <https://doi.org/10.1128/mBio.01862-15>.
 66. Ball B, Geddes-McAlister J. 2019. Quantitative proteomic profiling of *Cryptococcus neoformans*. *Curr Protoc Microbiol* 55:e94.
 67. Santiago-Tirado FH, Peng T, Yang M, Hang HC, Doering TL. 2015. A single protein S-acyl transferase acts through diverse substrates to determine cryptococcal morphology, stress tolerance, and pathogenic outcome. *PLoS Pathog* 11:e1004908. <https://doi.org/10.1371/journal.ppat.1004908>.
 68. Cadieux B, Lian T, Hu G, Wang J, Biondo C, Teti G, Liu V, Murphy M, Creagh AL, Kronstad JW. 2013. The Mannoprotein Cig1 supports iron acquisition from heme and virulence in the pathogenic fungus *Cryptococcus neoformans*. *J Infect Dis* 207:1339–1347. <https://doi.org/10.1093/infdis/jit029>.